

DINARIC KARST POLJES – FLOODS FOR LIFE

Edited by: Peter Sackl,
Romy Durst, Dražen Kotrošan
and Borut Stumberger

euRONATUR FOUNDATION



DINARIC KARST POLJES – FLOODS FOR LIFE

**Edited by: Peter Sackl,
Romy Durst, Dražen Kotrošan
and Borut Stumberger**

Proceedings of the 1st Workshop on Karst Poljes as
Wetlands of National and International Importance,
Livno, 30 September - 1 October 2014

euRONATUR FOUNDATION



Olm *Proteus anguinus* (Photo: Gregor Aljančič)

ENDANGERED PROTEUS: COMBINING DNA AND GIS ANALYSES FOR ITS CONSERVATION

Gregor Aljančič¹, Špela Gorički¹, Magdalena Năpăruș^{1,2}, David Stanković³ & Matjaž Kuntner⁴

¹Society for Cave Biology, Tular Cave Laboratory, Oldhamska c. 8a, SI-4000 Kranj, Slovenia, E-mail: gregor.aljancic@guest.arnes.si, goricki.spela@gmail.com

²LASIG, École Polytechnique Fédérale de Lausanne, Station 18, CH-1015 Lausanne, Switzerland, E-mail: magda.naparus@gmail.com

³University of Ljubljana, Biotechnical Faculty, Zootechnical Department, Groblje 3, SI-1230 Domžale, Slovenia, E-mail: david.l.stankovic@biologija.org

⁴Institute of Biology, Scientific Research Centre, Slovenian Academy of Sciences and Arts, Novi trg 2, SI-1000 Ljubljana, Slovenia, E-mail: kuntner@gmail.com

Summary

Proteus anguinus, an endemic amphibian of the subterranean waters of the Dinaric Karst, is restricted to its cave habitat. However, during seasonal flooding, some specimens are washed-out of their subterranean environment. While this may be considered as a highly hazardous way for *Proteus* to disperse into new habitats, it is obvious that all these individuals present a constant loss to their population. The Tular Cave Laboratory serves as a sanctuary for injured specimens accidentally washed-out of their subterranean habitat during seasonal flooding. Since 2008, 17 cases have been documented in Slovenia, and 7 of these animals were successfully returned to their source populations. Although the periodic loss of individual animals has been well balanced through the evolution of *Proteus*, a concern is raised when possible effects of climate change, large-scale hydrotechnical works and agriculture intensification are considered: the timing, frequency and magnitude of flood events are expected to be changed within a very short period of time. Here we discuss the risks and propose the actions necessary to halt the loss of these rare and highly endangered animals due to man-induced changes in flood regimes of the karst poljes.

Firstly, before any animal is returned to nature, the veterinary care and a strict protocol should minimize the transmission of potential infection. Secondly, if washed-out individuals are to be returned to nature, their source population must be accurately identified. Screening for DNA markers powerful enough to detect ongoing gene-flow, such as micro-satellites and single-nucleotide polymorphisms (SNPs), should minimize the potential for genetic pollution. Thirdly, the washed-out individuals deposited on karst fields often cannot be returned directly to their local cave system, since only small fragments of

Proteus subterranean habitat can be accessed by man.

We are currently developing a method of detection of traces of *Proteus* DNA in water samples (environmental DNA); when integrated in an accurate distribution Geographic Information System (GIS) model, the potential pattern of its genetic variability within the complex karst landscape will be determined. The resulted database should then be referenced not only to guide the return of washed-out individuals to nature but also to minimize the potential impact of any planned hydrotechnical and water-extraction activities in karst poljes on the genetic integrity of *Proteus* populations.

Sažetak

Proteus anguinus, endemični vodozemac podzemnih voda dinarskog krša, vezan je za pećinska staništa. No, tokom sezonskog plavljenja, neki primjerci budu izbačeni vodenom strujom iz svog podzemnog okruženja. Sa jedne strane, to se može smatrati veoma riskantnim načinom kojim *Proteus* naseljava nova staništa, ali je očito i da ovakve jedinke predstavljaju stalni gubitak za populaciju. Jamski laboratorij Tular služi kao utočište za povrijeđene primjerke koji su slučajno izbačeni iz svog podzemnog staništa tokom sezonskih poplava. Od 2008. u Sloveniji je zabilježeno 17 slučajeva, a 7 tih životinja je uspješno vraćeno u populacije iz kojih su potekle. Iako je povremeni gubitak jedinki nešto sa čime se ova vrsta sustetala tokom evolucije, postoji zabrinutost kada se u obzir uzmu moguće posljedice klimatskih promjena, hidrotehničkih radova i razvoja poljoprivrede: vrijeme, učestalost i veličina poplava će se vjerovatno promijeniti u veoma kratkom vremenskom periodu. U ovom radu bavimo se rizicima i predlažemo radnje koje su neophodne da se zaustavi gubitak ovih rijetkih i veoma ugroženih životinja zbog promjena u režimu poplava u kraškim poljima, koje je čovjek uzrokovao. Prvo,

prije nego što se ijedna životinja vrati u prirodu, veterinarska njega i strogi protokol treba smanjiti prenos moguće infekcije. Drugo, ako se izbačene jedinke trebaju vratiti u prirodu, treba tačno odrediti iz koje populacije su potekle. Screening DNA markera koji su dovoljno jaki da otkriju protok gena, kao što su mikrosateliti i jednonukleotidni polimorfizmi (SNPs) trebali bi smanjiti mogućnost genetičkog onečišćenja. Treće, jedinke koje su izbačene iz podzemnih staništa i nađene na kraškim poljima se obično ne mogu direktno vratiti u lokalni pećinski sistem, jer je samo malidio podzemnih staništa ove vrste dostupan čovjeku. Trenutno razvijamo metodu za otkrivanje tragova DNA *Proteusa* u uzorcima vode (okolišna DNA); kada se to ujedini sa stvarnim GIS (Geografski Informacioni Sistem) modelom rasprostranjenja, potencijalni režim genetske raznolikosti unutar kompleksa kraških polja će biti određen. Baza podataka koja će biti rezultat toga bit će korištena ne samo kao vodič za vraćanje izbačenih jedinki u prirodu, nego i za smanjenje potencijalnog uticaja planiranih hidrotehničkih aktivnosti u kraškim poljima na genetički integritet populacija *Proteusa*.

Keywords: *Proteus*, Amphibia, environmental DNA, karst ecohydrology, vulnerability map

Introduction

The Dinaric Karst is one of the world's prime heritage landscapes, both for natural and cultural phenomena. Dinaric Karst is also important for the study of evolutionary and ecological interplay. For example, here the endemic subterranean vertebrate fauna meets migrating birds, both animal groups having been affected by the single geological evolution of karst polje and its periodical flooding (sensu Bonacci *et al.* 2008), and subject to natural selection of over ten million years (Trontelj *et al.* 2007). The flagship species of this unique natural diversity is the olm, *Proteus anguinus* Laurenti 1768, a true symbol of karst and its history of research (M. Aljančič *et al.* 1993). Periodical flooding has probably been an important selective force in the evolution of this cave-dwelling amphibian (Aljančič & Năpăruș 2012), as well as of some populations of endemic surface-dwelling fish (*Telestes*, *Phoxinellus* and *Delminichthys*) (see also Palandačić *et al.* 2012). Periodical flooding in this karst system, however, also influences the occurrence of migratory birds of the Adriatic Flyway; these represent transitory, yet punctual guests that are also synchronized with periodical flooding of the Dinaric Karst

(Schneider-Jacoby *et al.* 2006; Stumberger 2010).

Through the centuries, human populations have also adapted to the karst landscape developing their own sustainable strategies. Recently, however, anthropogenic activities are largely negative, coming from intensive agricultural activities (e.g., Slovenia: overuse of biogas slurry fertilizer/ B. Bulog & A. Hudoklin, pers. comm.; Bosnia and Herzegovina: converting pastures into arable land/ B. Stumberger, pers. comm.), energy production (hydro and thermal power plants), and unregulated urbanization. Southeast Herzegovina, in particular, has seen these human activities reaching catastrophic proportions (see the case of Popovo Polje; M. Aljančič 1963, Čučković 1983, Lewarne 1999, Lučić 2013).

The case of *Proteus*: floods as a constant threat?

Flooding—an important periodic event in caves—is a transport agent for organic matter (also for contaminants) as well as for organisms, affecting their colonization, dispersal, and life cycles. In common with all stygobionts, *Proteus* is entirely restricted to its cave environment, and no longer able to be ecologically competitive in surface habitats. However, during flooding, a number of specimens get washed-out onto the surface, where they become stranded away from their subterranean environment and become exposed to predation on the open surface of the karst polje. It is reasonable to speculate that the earliest human knowledge of the

While becoming stranded on the surface may in fact be a part of natural history of *Proteus* intended to enable dispersal into new habitats, this highly hazardous strategy is, obviously, very costly in terms of loss of individuals from source populations.

existence of subterranean fauna was through encounters with such washed-out animals. As early as 1689 J. V. Valvasor encountered *Proteus* precisely under these circumstances, and provided its first description.



Fig. 1: Proteus stranded in Kljunov ribnik near Pivka, Slovenia, 29 December 2008; serious chilblains on dorsal side (Photo: Gregor Aljančič)

While becoming stranded on the surface may in fact be a part of natural history of *Proteus* intended to enable dispersal into new habitats, this highly hazardous strategy is, obviously, very costly in terms of loss of individuals from source populations. The fate of stranded individuals is quite predictable as the odds to re-enter the underground and thus to survive are minimal. The animals are often deposited on temporarily flooded fields and may survive for up to several weeks and as long as high waters persist; they may suffer sunburns or chilblains, and eventually desiccate (Fig. 1). Others may be carried further into surface streams where they get preyed upon by fish, birds or other predators.

We presume that *Proteus* has developed several responses to reduce the danger of being washed out of its subterranean habitat (Aljančič & Prelovšek 2010) as well as adaptations of its feeding and reproductive behaviour. Due to the extremely long lifespan on the one hand (animals may survive nearly 100 years in captivity) and long reproduction cycles on the other (*Proteus* reproduce approximately every 8 years in captivity), each individual loss may be fairly detrimental for the size of its population. Although this periodic loss has presumably been an annual constant through its evolution, more rapid changes due to human induced climate change, which are expected to change the timing, frequency and magnitude of floods dramatically, will most probably exceed the ability of *Proteus* to cope with environmental changes.

The Tular Cave Laboratory has extensively studied this neglected phenomenon since 2008, documenting 17 cases

of stranded *Proteus* in Slovenia. All animals were found by chance and reported by locals. Through this research we unexpectedly became involved in a rescue mission: seven of these animals were saved and returned to their source population (Aljančič & Năpăruș 2012), while injured animals found sanctuary in the Tular Cave Laboratory.

Why location matters

When an individual found on the surface cannot be returned directly to the cave or spring from which it was washed out by flood, the data on the exact distribution of its source population can guide its return. Analyses of mitochondrial DNA sequences revealed distinct lineages of *Proteus* from the Dinaric Karst of (1) Istria, (2) Dalmatia-Herzegovina, (3) Bosanska Krajina, (4) Lika, (5) SW Slovenia and (6) SE Slovenia (Gorički 2006; cf. Fig. 4 in Gorički & Trontelj 2006; Trontelj *et al.* 2007, 2009; and unpublished data). A study of highly variable nuclear DNA sequences is still lacking, but will, hopefully, resolve the complex network of relationships on a more local scale, i. e. within the populations that mitochondrial DNA indicated. To clearly delimit populations with on-going gene flow, intensive sampling coupled with datasets of karst features and their spatial relationships within the karst system, datasets with geological and hydrogeological features and their spatial relationships, a high resolution digital elevation model of the area, and spatial statistical modelling are needed. This is especially significant in the case of populations with very small ranges or which inhabit areas that have received little attention in the past.

The Tular Cave Laboratory is developing an indirect, forensic approach (which will be published elsewhere) to facilitate the search for *Proteus* in even the most inaccessible locations. Namely, during the process of skin growth and regeneration, fragments of epidermal cells, along with the DNA they contain, are constantly shed from the skin of aquatic vertebrates and carried away by water. Such DNA dissolved in water is called environmental DNA (eDNA). In its most basic form, the methodology aims at detecting traces of *Proteus* DNA released in water; as new genetic markers are found, it can be expanded for fast and routine genotypization of water samples. DNA extracted from the mucous of a washed-out individual is then compared to the integrated DNA-GIS database. In this way it will be possible to determine the origin of the washed-out animal, and the appropriate site to release the animal can be identified.

Further implications of combined DNA and GIS analysis for *Proteus* conservation

Due to their high specialization to a narrow range of abiotic conditions in the subterranean environment, all groundwater organisms are extremely vulnerable to direct and indirect alterations of their habitats. In Southeastern Bosnia and Herzegovina, large scale hydrotechnical activities are the main cause for disturbances to the natural flooding regime of caves and karst poljes, reduction of the catchment area, land amelioration, and water pumping (Ozimec 2011). Through reduction or loss of aquatic habitats in caves, or microclimatic and ecological changes such as temperature increase, reduction of dissolved oxygen, and reduction or increase in the quantity of organic matter (Ozimec 2011) these activities may have a detrimental impact on the density of groundwater fauna, including its flagship species *Proteus* (e. g., the case of Popovo Polje; M. Aljančič 1963, Čučković 1983, Lewarne 1999, Lučić 2013). In particular, intensive engineering works to divert the waters from Dabarsko polje towards Fatničko polje and further towards Bileća, Trebinje, and the Ombla hydroelectric plant near Dubrovnik considerably reduce the flow of groundwater towards the lower Neretva River and its delta (J. Mulaomerović, pers. comm.). Furthermore, a substantial portion of waters from Popovo polje is being diverted towards Ombla spring, while only a small part is directed to the reversible hydroelectric plant “Čapljinina”, located on the eastern side of the Svitavsko-Deransko polje. The reduction of input of groundwater from karst poljes of Eastern Herzegovina into the Neretva River’s lower course and the delta results in an increase of saltwater penetrating further upstream. The observed increase of salinity may have a direct adverse effect on the localities of *Proteus* in the Neretva River delta and Hutovo Blato. Changes in the direction of watercourses may also affect the gene pool of *Proteus* populations (Sket 1997). Existing legal acts – with Bosnia and Herzegovina being the only political entity within the range of *Proteus* that lacks any legislation enforcing its conservation – do not entirely protect aquatic cave animals and their habitats from negative human impacts on groundwater integrity. A decline of several populations of *Proteus* has been reported, and in some localities *Proteus* has already become extinct (Sket 1997). The IUCN Red List of Threatened Species defines *Proteus anguinus* as vulnerable, and recommends urgent measures to revert its population decline (Arntzen *et al.* 2009). The extent of its decline, however, cannot be

estimated without an extensive survey of its distribution. Current knowledge on the distribution of *Proteus* in Bosnia and Herzegovina (Kotrošan 2002) is particularly scarce in the area along the lower course of the Neretva Rivier and its tributaries. In the area of Hutovo blato, only a single *Proteus* locality is known: a well in the village Čore near the Babino oko spring. Undoubtedly, *Proteus* can also inhabit other caves in the area. This is supported by the oldest known depiction of *Proteus* in Bosnia and Herzegovina, on a ‘stećak’ (monumental gravestone typical for Medieval Bosnia) from ca. 1477, found in the Boljuni necropolis near Stolac (Mulaomerović & Hodžić 2012), which conveys *Proteus* as having a long symbolic presence.

Conclusions

The Tular Cave Laboratory has started utilizing the eDNA to infer the presence of *Proteus* in the most threatened sites. The results of this intensive survey will provide a scientific basis for enforcing the long-term protection of *Proteus* populations and its habitats, and will help to mitigate current and future threats. The highly efficient, non-invasive, DNA-based method to detect *Proteus* from water samples coupled with a set of spatial data will provide a vulnerability map of *Proteus* which will visualize zones most threatened by human impacts along with the most active threats. When integrated in an accurate Geographic Information System (GIS) distribution model, the patterns of the genetic variability of *Proteus* within the complex karst landscape will emerge. The DNA-GIS database will further provide the scientific basis for the return of washed-

The highly vulnerable *Proteus* has a huge potential to become a symbol of a successful balance between conservation and sustainable management of the karst environment.

out individuals to their source cave system. Furthermore, it will also help to prepare urgently needed vulnerability models for assessing potential impacts of hydrotechnical and water-extraction activities in karst poljes on the

genetic structure of *Proteus* populations. The accuracy of the resulting distribution and vulnerability models will be validated by spatial statistical analysis (Năpăruș & Kuntner 2012; Meleg et. al 2013). This analytical tool is designed to implement future conservation action plans needed for building sustainable strategies for landscape management and ecological forecasting. Future activities may even reach beyond local geography and specifics of the distribution of *Proteus* in the Dinaric Karst – with the present survey serving as a model for future assessments of the vulnerability of aquatic cave fauna worldwide – and, thus, offer the possibility to implement science based conservation strategies for a sustainable management of the karst environment and its biodiversity.

The present study has strengthened our belief that a complex ecosystem of Dinaric Karst poljes (above and underground) can only be preserved through multidisciplinary scientific efforts and with the support of international nature conservation organizations. An active alliance of organisations committed to nature conservation of the Dinaric Karst is urgently needed in order to embark on an ambitious task to promote, share and implement an advanced action plan to save *Proteus*, Europe's only cave vertebrate, along its fragmented range in the Western Balkans. The highly vulnerable *Proteus* has a huge potential to become a symbol of a successful balance between conservation and sustainable management of the karst environment.

Acknowledgements

We would like to thank Borut Stumberger and Peter Sackl for helpful comments on the manuscript, and Brian Lewarne for linguistic review. We thank Ivan Bebek, Ilhan Dervović, Jelena Kadić, Dražen Kotrošan, Brian Lewarne, Ivo Lučić, Jasminko Mulaomerović, Dušan Musa, Goran Panić and Sara Todorović for the valuable information on greatest threats to *Proteus* in Bosnia and Herzegovina. We also thank EuroNatur and Naše ptice for hosting the workshop. The eDNA survey in Southeastern Herzegovina and Montenegro is performed in partnership with the Center for Karst and Speleology (Sarajevo), the Republic Institute for Protection of Cultural, Historical and Natural Heritage (Banja Luka), the Biospeleological Society of Montenegro (Podgorica), the Herzegovinian Mountain Rescue Service (Mostar), and with the financial support of the Critical Ecosystem Partnership Fund/BirdLife International and DOPPS, the University of Maryland, and the Institute of Biology at the Scientific Research Centre (Ljubljana).

References

- Aljančič G., Prelovšek M. (2010):** Does *Proteus* detect and react to a sudden rise of water conductivity which indicates incoming flood? Abstract Book 20th International Conference on Subterranean Biology. Postojna, Slovenia; pp. 114 – 115.
- Aljančič G., Năpăruș M. (2012):** Stygobionts washed out to surface: A case of *Proteus anguinus*. Abstract Book 21st International Conference on Subterranean Biology. Košice, Slovakia; pp. 22 – 23.
- Aljančič M. (1963):** Po jamah Popovega polja [In caves of Popovo Polje]. *Proteus* 25(9-10): 239 – 245.
- Aljančič M., Bulog B., Kranjc A., Josipovič D., Sket B., Skoberne P. (1993):** *Proteus*: The Mysterious Ruler of Karst Darkness. Vitrum, Ljubljana, 75 pp.
- Arntzen J.W., Denoël M., Miaud C., Andreone F., Vogrin M., Edgar P., Crnobrnja Isailović J., Ajtic R., Corti C. (2009):** *Proteus anguinus*. In: IUCN (2013), IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>. Downloaded on 25 November 2013.
- Bonacci O., Pipan T., Culver D. C. (2008):** A framework for karst ecohydrology. *Environmental Geology* 65(5): 891 – 900.
- Čučković S. (1983):** Uticaj promjene režima vodotoka hidrosistema Trebišnjica na faunu kraškog podzemlja [The influence of the change in the water-course regime of the Trebišnjica water-system on the fauna of the karst underground regions]. *Naš Krš* 9: 129 – 142.
- Gorički Š. (2006):** Filogeografska in morfološka analiza populacij močerila (*Proteus anguinus*). [Phylogeographic and morphological analysis of European cave salamander (*Proteus anguinus*) populations.] Ph.D. Dissertation, Ljubljana, Slovenia, 76 pp.
- Gorički Š., Trontelj P. (2006):** Structure and evolution of the mitochondrial control region and flanking sequences in the European cave salamander *Proteus anguinus*. *Gene* 378: 31 – 41.
- Kotrošan D. (2002):** Rasprostranjenje čovječije ribice na području Bosne i Hercegovine [Distribution of the *Proteus anguinus* Laurenti, 1768] in Bosnia and Herzegovina]. *Naš Krš* 35: 57 – 64.
- Lewarne B. (1999):** A joint strategy for the protection of the Trebinje *Proteus anguinus* and its natural karst habitat. *Naš Krš* 32: 35 – 50.
- Lučić I. (2013):** Obrnuti tokove vode i povijesti [Turning course of water and history]. *Međunarodni forum Bosne – IFB*, 61/62(13): 75 – 116.
- Meleg I. N., Năpăruș M., Fiers F., Meleg H. I., Vlaicu M., Moldovan O. T. (2013):** The relationships between land cover, climate and cave copepod spatial distribution and suitability along the Carpathians. *Environmental Conservation*. FirstView: 1-11.
- Mulaomerović J., Hodžić M. (2012):** *Proteus anguinus* in Bosnia and Herzegovina: from the Middle ages to today. Abstracts 21st International Conference on Subterranean Biology, Košice, Slovakia; p. 77.
- Năpăruș M., Kuntner M. (2012):** A GIS Model Predicting Potential Distributions of a Lineage: A Test Case on Hermit Spiders (Nephilidae: Nephilengys). *PLoS ONE* 7/1: e30047.doi:10.1371/journal.pone.0030047.
- Ozimec R. (2011):** Red book of Dinaric cave fauna – an example from Croatia. In: Prelovšek M., Zupan Hajna N. (eds.), *Pressures and Protection of the Underground Karst: Cases from Slovenia and Croatia*. Karst Research Institute ZRC SAZU, Postojna; pp. 182 – 190.
- Palandačić A., Bonacci O., Snoj A. (2012):** Molecular data as a possible tool for tracing groundwater flow in karst environment: example of *Delminichthys adspersus* in Dinaric karst system. *Ecohydrology* 5/ 6: 791 – 797.
- Schneider-Jacoby M., Rubinić B., Sackl P., Stumberger B. (2006):** A preliminary assessment of the ornithological importance of Livanjsko Polje (Cetina River Basin, Bosnia and Herzegovina). *Acrocephalus* 27: 45 – 57.
- Sket B. (1997):** Distribution of *Proteus* (Amphibia: Urodela: Proteidae) and its possible explanation. *Journal of Biogeography* 24: 263 – 280.
- Stumberger B. (2010):** A classification of karst poljes in the Dinarides and their significance for waterbird conservation. In: Denac D., Schneider-Jacoby M., Stumberger B. (eds.), *Adriatic Flyway – Closing the Gap in Bird Conservation*. EuroNatur, Radolfzell; pp. 69 – 78.
- Trontelj P., Gorički Š., Polak S., Verovnik R., Zakšek V., Sket B. (2007):** Age estimates for some subterranean taxa and lineages in the Dinaric Karst. *Acta carsologica* 36: 183 – 189.
- Trontelj P., Douady C. J., Fišer C., Gibert J., Gorički Š., Lefébure T., Sket B., Zakšek V. (2009):** A molecular test for cryptic diversity in ground water: how large are the ranges of macro-stygobionts? *Freshwater Biology* 54: 727 – 744.